Scaling Crossover Description of Near-Critical Vapor-Liquid Equilibria in Aqueous Solutions of Sodium Chloride

A.A. Povodyrev, M.A. Anisimov, and J.V. Sengers
Institute for Physical Science and Technology
University of Maryland
College Park, MD 20742 USA

J.M.H. Levelt Sengers

Physical and Chemical Properties Division

National Institute of Standards and Technology
Gaithersburg, MD 20899 USA

An adequate interpretation of experimental data in binary fluids near the vapor-liquid critical point is complicated by an interplay between two different crossover phenomena: crossover between pure-fluid-like behavior and mixture-like behavior; and crossover from Ising-like asymptotic scaling behavior to mean-field behavior. Moreover, in ionic systems possible dissociation-association processes cause additional complications. We present a quantitative description of experimental data for vapor-liquid equilibria, critical locus, and isochoric specific heat capacity of near-critical aqueous solutions of sodium chloride in terms of a crossover model based on the extended principle of critical-point universality. We have analyzed existing experimental data on coexisting concentrations and densities along isotherm (above the critical point of water) in a wide range of pressures (from the critical pressures $P_{\rm c}$ to $P/P_{\rm c} \approx 0.5$), and isochoric specific heat capacity data along critical isochores. It is shown that the critical phenomena in this system are characterized by a crossover from Ising-like asymptotic behavior to mean-field behavior and by a decrease of the Ising-like asymptotic critical region with increase of sodium chloride overall concentration. However, the model is inadequate at the lowest overall concentrations that may be due to dissociation-association effects.

The research of A.A. Povodyrev is supported by the American Chemical Society under Petroleum Research Fund Grant 30693-AC9. The research of M.A. Anisimov and J.V. Sengers is supported by DOE Grant FG02-95ER-14509.